



GOVERNMENT OF INDIA
Ministry of Agriculture & Farmers Welfare
Department of Agriculture, Cooperation & Farmers Welfare
Directorate of Plant Protection, Quarantine & Storage
Central Insecticides Board & Registration Committee
N.H. IV, Faridabad-121 001

MAJOR USES OF PESTICIDES

Registered under the Insecticides Act, 1968

30.06.2016

Disclaimer: The document has been compiled on the basis of available information for guidance and not for legal purposes.

HERBICIDES

1. Herbicides products approved uses: Page 1 to 32
2. Herbicides combinations approved uses: Page 33 to 39

APPROVED USES OF REGISTERED HERBICIDES

HERBICIDES

Herbicide name & approved Crops	Weed species	Dosage /ha		Dilution In Water (Litres)	Waiting period / PHI between last application & harvest (days)
		a.i. (gm/ Kg)	Formulation in (gm/ml /Kg/ ltr)		
Alachlor 50% EC					
Cotton	<i>Digera arvensis</i> <i>Echinochloa colonum</i> , <i>Eragrostis major</i> <i>Euphorbia hirta</i> <i>Phyllanthus niruri</i> <i>Portulaca oleracea</i> <i>Trianthema portulacastrum</i> <i>Flaveria australasica</i> <i>Gynandropsis pentaphylla</i>	2-2.5 kg	4-5 ltrs.	250-500	210-240
Maize	<i>Echinochloa colonum</i> , <i>Euphorbia hirta</i> <i>Eleusine indica</i>	2.5 kg	5 ltrs.	250-500	90
Groundnut	<i>Amaranthus viridis</i> <i>Digitaria spp.</i> <i>Echinochloa spp.</i> <i>Euphorbia hirta</i> <i>Phyllanthus niruri</i> <i>Portulaca oleracea</i> <i>Trianthema portulacastrum</i>	2.5 kg	5 ltrs.	250-500	120-150
	<i>Acanthosermum hispidum</i> <i>Flaveria australasica</i>	1.5-2.5 kg	3-5 ltrs.	250-500	120-150

Soybean	<i>Amaranthus viridis</i> <i>Cleome viscosa</i> <i>Cyperus iria</i> <i>Dactyloctenium aegyptium</i> <i>Echinochloa spp.</i> <i>Eleusine indica</i> <i>Setaria glauca</i>	2.5 kg	5 ltrs.	250-500	
Alachlor 10% GR					
Cotton	<i>Dactyloctenium aegyptium</i>	2.0-2.5 Kg	20-25 Kg	-	-
Maize / Groundnut / Soybean	<i>Digitaria spp.,</i> <i>Echinochloa spp.,</i> <i>Chenopodium album</i>	1.5-2.5 Kg	15-25 Kg	-	-
Anilofos 30% EC					
Transplanted paddy	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis,</i> <i>Cyperus iria,</i> <i>Eclipta alba</i> <i>Ischaemum rugosum</i> <i>Fimbristylis sp.</i> <i>Marsilea quadrifoliata</i>	0.3-0.45 kg	1-1.5 ltrs.	375-500	30
Anilofos 18% EC					
Transplanted Paddy	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis,</i> <i>Cyperus iria,</i> <i>Eclipta alba</i> <i>Ischaemum rugosum</i> <i>Fimbristylis sp.</i>	0.30-0.45 kg	1.66-2.5 kg	500-600	-

Anilophos 2 % G					
Transplanted rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Ischaemum rugosum</i> <i>Cyperus iria</i> , <i>Cyperus difformis</i> , <i>Fimbristylis sp.</i>	0.4-0.5 Kg	20-25 Kg	-	30
Atrazine 50% WP					
Maize	<i>Trianthema monogyna</i> <i>Digera arvensis</i> , <i>Echinochloa spp</i> <i>Eleusine Spp.</i> <i>Xanthium strumarium</i> <i>Brachiaria sp, Digitaria</i> <i>sp, Amaranthus viridis</i> , <i>Cleome viscosa</i> , <i>Polygonum spp.</i>	0.5-1.0 kg	1-2 kg	500-700	-
Azimsulfuron 50% DF					
Rice (Transplanted)	<i>Enchinochloa colonum</i> , <i>E. crusgalli</i> , <i>Cyperus spp.</i> , <i>Fimbristylis miliacea</i> , <i>Ludwigia parviflora</i> , <i>Eclipta alba</i> , <i>Bergia</i> <i>capensis</i> , <i>Marsilea</i> <i>quadrifoliata</i> , <i>Ammania</i> <i>baccifera</i> , <i>Sphenoclea</i> <i>zeylanica</i>	35	70	300	59
Rice (Direct Seeded)	<i>Enchinochloa colonum</i> , <i>E. crusgalli</i> , <i>Cyperus spp.</i> , <i>Fimbristylis miliacea</i> , <i>Ludwigia parviflora</i> , <i>Eclipta alba</i> , <i>Bergia</i> <i>capensis</i> , <i>Marsilea</i> <i>quadrifoliata</i> , <i>Ammania</i> <i>baccifera</i> , <i>Sphenoclea</i> <i>zeylanica</i>	35	70	300	59

Bensulfuron Methyl 60% DF					
Transplanted Rice. Pre-em (3 DAT)	<i>Marsilea quadrifoliata</i> <i>Eclipta alba</i> , <i>Ammania baccifera</i> , <i>Ludwigia parviflora</i> , <i>Sphenoclea Zeylenica</i> , <i>Monochoria vaginalis</i> , <i>Alternanthera sessillis</i> <i>Cyperus iria</i> , <i>Cyperus difformis</i> , <i>Fimbristylis miliacea</i> , <i>Scirpus royeli</i>	60 gm	100 gm	300 ltrs	88 days
Transplated Rice (post-em 20 DAT)	<i>Ammania baccifera</i> <i>Cyperus differmis</i> <i>Cyperus iria</i> <i>Eclipta alba</i> <i>Fimbristylis miliacca</i> <i>Ludwigia parviflora</i> <i>Marsilea quadrifoliata</i> <i>Monochoria vaginalis</i> <i>Alternanthera sessillis</i> <i>Scirpus royeli</i> <i>Sphenoclea zeylenica</i>	60gm	100 gm	300 ltrs.	71
Bentazone 480 g/l SL					
Soybean	<i>Acalypha indica</i> <i>Commelina benghalensis</i> , <i>Cyperus rotundus</i> <i>Echinochloa colonum</i> <i>Echinochloa crusgalli</i>	960	2000	500	62
Transplated Rice	<i>Cyperus differmis</i> <i>Cyperus rotundus</i> <i>Echinochloa colonum</i> <i>Echinochloa crusgalli</i> <i>Eclipta alba</i> <i>Ludwigia parviflora</i>	960	2000	500	71

Bispyribac Sodium 10% SC					
Rice (Nursary)	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i>	20 gm	200 ml.	300 ltrs.	-
Rice (Transplanted)	<i>Ischaemum rugosum</i> <i>Cyperus difformis</i> , <i>Cyperus iria</i> ,	20 gm	200 ml	300 ltrs.	78
Rice (Direct seeded)	<i>Fimbristylis miliacea</i> , <i>Eclipta alba</i> , <i>Ludwigia parviflora</i> , <i>Monochoria vaginalis</i> , <i>Alternanthera philoxeroides</i> , <i>Sphenoclea zeylanica</i>	20 gm	200 ml	300 ltrs.	78
Butachlor 50% EC					
Paddy (transplanted)	<i>Cyperus difformis</i> <i>Cyperus iria</i> <i>Echinochloa crusgalli</i> , <i>Echinochloa colonum</i> , <i>Eleusine indica</i> , <i>Eclipta alba</i> , <i>Fimbristylis miliacea</i> , <i>Ludwigia parviflora</i> , <i>Sphenoclea zeylanica</i>	1.25- 2.00kg	2.5-4 ltrs	250-500	90-120
Butachlor 5% GR					
Rice	<i>Cyperus difformis</i> <i>Cyperus iria</i> , <i>Echinochloa crusgalli</i> , <i>Echinochloa colonum</i> , <i>Eleusine indica</i> , <i>Eclipta alba</i> , <i>Fimbristylis miliacea</i> , <i>Ludwigia purviflora</i> , <i>Sphenoclea zeylanica</i>	1.25-2.0 kg	25-40kg.	-	90-120
Butachlor 50 % EW					

Transplanted Rice	<i>Echinochloa colonum</i> <i>Echinochloa crusgalli</i> , <i>Cyperus difformis</i> <i>Cyperus iria</i> <i>Eclipta alba</i> , <i>Fimbristylis miliacea</i> <i>Ludwigia parviflora</i> , <i>Sphenoclea zeylanica</i> <i>Monochoria vaginalis</i>	1.25-1.5 Kg	2.5-3.0	2.50-500	-
Carfentrazone ethyl 40% DF					
Wheat	<i>Chenopodium album</i> , <i>Melilotus Indica</i> , <i>Melilotus alba</i> , <i>Medicago denticulata</i> , <i>Lathyrus aphaca</i> , <i>Analgalis arvensis</i> , <i>Vicia sativa</i> <i>Circium</i> <i>arvense</i> , <i>Rumex sp</i> , <i>Malwa sp.</i>	20gm	50 gm.	400	80
Chlorimuron Ethyl 25% WP					
Soybean	<i>Eclipta alba</i> , <i>Commelina</i> <i>benghalensis</i> <i>Chenopodium album</i> , <i>Cyperus rotundus</i> , <i>Cyprus iria</i> , <i>Parthenium</i> <i>hysterophorus</i> , <i>Acalypha indica</i> , <i>Phyllanthus niruri</i> , <i>Trianthema</i> <i>portulacashurm</i> , <i>Caesulia auxillaris</i>	9 gm	36 gm.	300 ltrs. +Surfact ant 0.2 % (Iso- octyl phenoxy- poloxeth anol 12.5 %)	45
Rice (transplanted)	<i>Echinochloa crusgalli</i> , <i>Eclipta alba</i> , <i>Commelina</i> <i>benghalensis</i> , <i>Chenopodium album</i> , <i>Cyperus rotundus</i> , <i>Echinochloa colonum</i>	6gm	24 gm.	500-600	60
Cinmethylin 10% EC					

Transplanted Rice	<i>Cyperus iria</i> <i>Fimbristylis milacea</i> <i>Monochoria vaginalis</i> <i>Commelina</i> <i>Benghalensis</i> <i>Echinochloa crusgalli</i> <i>Marsilea minuta</i>	75-100 gm	0.75-1.0 ltrs.	500-700	110
Clodinafop- propargyl 15%WP					
Wheat	<i>Phalaris minor</i> (<i>Canary grass</i>)	60gm	400 gm.	375-400	110
Clomazone 50%EC					
Soybean	<i>Digiteria sp.</i> <i>Echinochloa sp.</i> <i>Parthenium</i> <i>hysterophorus</i> <i>Commelina sp.</i>	0.75- 1.00Kg	1.5-2.0 Ltrs.	500-600	90
Transplanted Rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis</i> <i>Cyperus iria, Ludwigia</i> <i>parviflora, Eclipta alba</i>	0.4 - 0.5kg	0.8-1.0 ltr	500-750	90
Cyhalofop Butyl 10% EC					
Rice (Directed seeded)	(<i>Echinochloa spp.</i>) <i>Barnyard grass</i>	75-80 gm	0.75- 0.80ltr	500-600	90
2,4-D Dimethyl Amine salt 58% SL					
Maize	<i>Trianthema monogyna,</i> <i>Amaranthus sp.,</i> <i>Tribulus terrestris,</i> <i>Boerhaavia diffusa,</i> <i>Euphorbia hirta,</i> <i>Portulaca oleracea,</i> <i>Cyperus sp.</i>	0.5 kg	0.86	400-500	50-60

Wheat	<i>Chenopodium album</i> , <i>Fumaria parviflora</i> , <i>Melilotus alba</i> , <i>Vicia</i> <i>sative</i> , <i>Asphodelus</i> <i>tenuifolius</i> , <i>Convolvulus arvensis</i> ;	0.5-0.75 kg	0.86-1.29	500-600	-
Sorghum	<i>Cyperus iria</i> , <i>Digera</i> <i>arvensis</i> , <i>Convolvulus</i> <i>arvensis</i> , <i>Trianthema</i> <i>sp.</i> , <i>Tridax</i> <i>procumbens</i> , <i>Euphorbia</i> <i>hirta</i> , <i>Phyllanthus</i> <i>niruri</i> .	1.8 kg	3.1	500-600	-
Potato	<i>Chenopodium album</i> , <i>Asphodelus tenuifolius</i> , <i>Anagalis arvensis</i> , <i>Convolvulus</i> <i>arvensis</i> , <i>Cyperus iria</i> , <i>Portulaca oleracea</i> .	2.0 kg	3.44	400	-
Sugarcane	<i>Cyperus iria</i> <i>Digitaria sp.</i> <i>Dactylactenium</i> <i>aegyptium</i> <i>Digera arvensis</i> <i>Portulaca oleracea</i> <i>Commelina</i> <i>benghalensis</i> <i>Convolvulus arvensis</i>	3.5	6.3	500	-
Aquatic Weeds	<i>Eichhornia crassipes</i> .	0.5-1.0 kg	0.86-1.72	600-700	15-20
Non crop area	<i>Parthenium</i> <i>hysterophorus</i> ,	2.65 kg	4.56	300-400	15-20
	<i>Cyperus rotundus</i>	2.5 kg	4.30	300-400	-
2,4-D Sodium salt Technical (having 2,4-D acid 80 % w/w) (Earlier Registered as 80%WP)					

Citrus	<i>Euphorbia spp.</i> <i>Convolvulus arvensis</i> <i>Coronopus didymus</i> <i>Amaranthus viridis</i> <i>Oxalis corniculata</i> <i>Tribulus terrestris</i> <i>Fumaria parviflora</i> <i>Sonchus arvensis</i>	1.00-2.5 kg	1.25-3.2 kg	600	>6 months
Grapes	<i>Convolvulus spp.</i> <i>Tridax procumbens</i>	2.0	2.5	500	> 90 days
Maize	<i>Amaranthus viridis,</i> <i>Trianthema portulacastrum</i> <i>Phyllanthus niruri.,Euphorbia geniculata,Amaranthus spinosus.Cleome chelidonii, Lagasca mollis</i>	1.00 Kg.	1.25	500	120(Pre-em) 90(post-em)
Sugarcane	<i>Boerhaavia diffusa</i> <i>Chenopodium album</i> <i>Tribulus terrestris</i> <i>Portulaca oleracea</i> <i>Xanthium spp.</i> <i>Convolvulus arvensis</i> <i>Amaranthus spinosus</i> <i>Digera arvensis</i> <i>Celosia argentina.</i>	2.0-2.6	2.5-3.25	600-900	300
Wheat	<i>Leucas aspera ,</i> <i>Chenopodium album,Vicia sativa, ,Argemone maxicana , Fimbristylis miliacea, Anagalis arvensis, Amaranthus spinosus.</i>	0.5-0.84 kg.	0.625-1.0	500	90
Aquatic Weeds	<i>Boerhaavia hispada,</i> <i>Eichhornia crassipes.</i>	1.5 kg	1.85.	600-1000	-
Non crop land	<i>Parthenium hysterophorus,</i> <i>Cyperus rotundus,</i> <i>Solanum elaeagnifolium.</i>	2.5-6.0 kg. 4-8 Kg 1.8 kg	3.2-7.5 5-10 2.25	600-1000 500-600 500-600	- - -

2,4-D Ethyl Ester 38 % EC (having 2,4-D acid 34 % w/w)					
Maize	<i>Trianthema monogyna</i> , <i>Amaranthus sp.</i> , <i>Portulaca oleracea.</i> , <i>Tribulus terrestris</i> , <i>Boerhaavia diffusa</i> , <i>Euphorbia hirta</i> , <i>Cyperus sp.</i>	0.9 kg	2.65 ltr	400-450	50-60
Sorghum	<i>Cyperus iria</i> , <i>Striga sp.</i> <i>Digera arvensis</i> , <i>Convolvulus arvensis</i> , <i>Trianthema sp.</i> , <i>Tridax</i> <i>procumbens</i> , <i>Euphorbia hirta</i> , <i>Phyllanthus niruri.</i>	1.0 kg	2.94	425	-
Transplanted Paddy	<i>Echinochloa colonum</i> , <i>Echinochloa crusgalli.</i>	0.85 kg	2.5	400	-
Wheat	<i>Chenopodium album</i> , <i>Asphodelus</i> <i>tenuifolius</i> , <i>Fumaria</i> <i>parviflora</i> <i>Melilotus</i> <i>alba.</i> <i>Spergula arvensis</i>	0.45-0.75 kg	1.32-2.2	450-500	-
Sugarcane	<i>Cyperus iria</i> , <i>Digitaria</i> <i>sp.</i> , <i>Dactyloctenium</i> , <i>Aegyptiana</i> , <i>Digera</i> <i>arevensis</i> , <i>Portulaca</i> <i>oleracea</i> , <i>Commelina</i> <i>benghalensis</i> , <i>amaranthus sp.</i> , <i>Convolvulus arvensis</i>	1.2 to 1.8	3.53- 5.29	500	300-330
Aquatic Weeds	<i>Eichhornia crassipes</i>	2.5 kg	7.5	700- 1000	-
2,4-D Ethyl Ester 4.5 % GR (having 2,4-D acid 4 % w/w)					

Transplanted Rice	<i>Echinochloa Coloum</i> <i>E. Crusgalli</i> <i>Panium ischaemum</i> <i>Cynodon dactylon</i> (germinating) <i>Cyperus rotundus</i> (germinating) <i>Cyperus iria</i> <i>C. difformis</i> <i>Ludwigia parviflora</i> <i>Monochoria Vaginalis</i> <i>Marsilea quadrifoliata</i> <i>Cyanotis cucutata</i> <i>Eclipta alba</i> <i>Ammania baccifera</i>	1.0 kg	25 kg	-	-
Diclofop Methyl 28% EC					
Wheat	<i>Avena fatua, Phalaris minor</i>	0.7-1.0 kg	2.5-3.5 ltr	500	90
Diuron 80% WP					
Cotton	<i>Amaranthus spp,</i> <i>Chenopodium album,</i> <i>Convolvulus arvensis</i> <i>Setaria glauca,</i> <i>Digitaria sp, Portulaca</i> <i>oleracea, Xanthium</i> <i>strumerium, Anagallis</i> <i>arvensis, Asphodelus</i> <i>temifolius, Euphorbia</i> <i>sp, Visia sativa</i> <i>Paspalum conjugatum,</i>	0.75-1.5 kg	1-2.2Kg.	625	-
Banana	<i>Cyperus iria,</i> <i>Commelina</i> <i>benghalensis, Digitaria</i> <i>sp, Amaranthus</i> <i>spp, Dactyloctenium, Ch</i> <i>loris barbata, Eragrostis</i> <i>zeylenica,</i>	1.60 kg	2 kg.	625	-
Rubber	<i>Grasses & Non</i> <i>grasses</i>	1.6-3.2 kg	2-4kg.	625	-
	<i>Cyperus</i>	0.8 kg	1.0 kg.	600	-

Maize	<i>iria, Echinochloa spp, Digitaria spp, Chenopodium album, Eleusine sp, Amaranthus sp, Phyllanthus niruri</i>				
Citrus (sweet orange)	<i>Cyperus iria, Tribulus Terrestris, Digera arvensis, Commelina nudiflora, Cocumis trigonus..</i>	2-4.0kg	2.5-5.0kg	600	-
Sugarcane	<i>Cyperus iria, Portulaca racea, Echinochloa rusgalli, Cynotis spp, Amaranthus spp, Convonvulus spp, Digitaria spp.</i>	1.6-3.2kg	2.0-4.0 kg.	600	-
Grapes	<i>Cleome viscose, Chenopodium album. Cyperus iria, Euphorbia hirta, Alternanthera echinata, Amaranthus spp, Argemone maxicana, Ipomoea spp, Xanthium strumerium, Fumeria parviflora, Asphodelus tenuifolius, Medicago denticulata, Eleusine aegyptia.</i>	1.6kg	2.0 kg.	625	-
Diclosulam 84% WDG					
Soyabean	Cyperus spp, Commilena benghalensis, Euphorbia geniculata, Digera arvensis, Acylipha spp, Echinochlo colona	22-26gm	26.2-30.9 Time of application 0-3 DAS	500	60
Ethoxysulfuron 15% WDG					

Transplanted Rice.	<i>Fimbristylis miliacea</i> <i>Cyperus iria</i> , <i>Cyperus difformis</i> , <i>Scirpus sp.</i> , <i>Eclipta alba</i> , <i>Marsilea quadrifoliata</i> , <i>Ammania baccifera</i> , <i>Monochoria vaginalis</i> ,	12.5-15gm	83.3-100gm	500	110
Fenoxaprop-p-ethyl 9.3% w/w EC (9% w/v)					
Soybean	<i>Echinochloa colonum</i> , <i>Echinochloa crusgalli</i> , <i>Digitaria sp</i> , <i>Eleusine indica</i> , <i>Setaria sp</i> , <i>Brachiaria sp.</i>	100gm.	1111 ml. (15-20 DAS)	250-300	100
Rice (transplaned)	<i>Echinochloa crusgalli</i> , <i>Echinochloa colona</i>	56.25 gm	625 ml. (10-15 DAT)	300-375	70
Blackgram	<i>Echinochloa crusgalli</i> , <i>Echinochloa colona</i> <i>Digitaria sp.</i> <i>Dactyloctenium</i> <i>Aegyptium</i>	56.25-67.5 g	625- 750ml. (15-20 DAS)	375-500	43
Cotton	<i>Echinochloa sp.</i> <i>Eluesine indica</i> <i>Dactyloctenium</i> <i>Aegyptium</i> <i>Eragrostit minor</i>	67.5 g	750ml. (20 -25 DAS)	375-500	87
Fenoxaprop-p-ethyl 10% EC					
Wheat	<i>Phalaris minor</i>	100-120gm	1.0-1.20 kg.	250-300	110
Fenoxaprop-p-ethyl 6.7% w/w EC					
Rice (Transplanted & Direct Seeded)	<i>Echinochloa sp.</i>	56.6- 60.38g	812.5-875	375-500	61
Fluazifop-p-butyl 13.4% EC					

Soybean	<i>Echinochloa colonum</i> , <i>Echinolchloa crusgalli</i> , <i>Eleusine indica</i> , <i>Cyanodon dactylon</i> , <i>Dactyloctenium</i> <i>Aegyptium</i> , <i>Digitaria sp.</i> , <i>Setaria sp.</i>	125-250 g	1000- 2000	500	90
Flucetosulfuron 10% w/w WG					
Rice (Transplanted)	<i>Echinochloa colonum</i> <i>Echinolchloa crusgalli</i> <i>Digitaria sanguinalis</i> <i>Paspalum discichum</i> <i>Paspalum</i> <i>scrobitulatum</i> <i>Leersia hexandra</i> <i>Panicum repens</i> <i>Setaria glauca</i> <i>Dinebra retroflexa</i> <i>Cyprus difformis</i> <i>Cyprus iria</i> <i>Fimbristylis miliaceae</i> <i>Alternanthera</i> <i>philoxeroides</i> <i>Alternanthera sessilis</i> <i>Marsilea quadrifolia</i> <i>Ammania baccifera</i> <i>Eclipta alba</i> <i>Eclipta prostrate</i> <i>Monochoria vaginalis</i> <i>Lindernia ciliate</i> <i>Ludwigia parviflora</i> <i>Sphenoclea zeylanica</i> <i>Commelina diffusa</i> <i>Cyanotis axillaris</i>	25	250	500	90
Fluchloralin 45% EC					

Cotton	<i>Acanthospermum hispidum, Cleome viscosa, Datura sp. Trianthema monogyna Tridax procumbens, Cynodon dactylon (germinating) Amaranthus spp., Portulaca spp, Achyranthus aspera, Euphorbia hirta, Cenchrus catharticus, Digitaria sanguinalis, Eleusine sp, Panicum sp, Lagasca mollis, Gynandropsis pentaphylla, Achalypha indica</i>	0.9-1.2kg	2.0-2.68 ltrs.	500-800	180
Soybean	<i>Eragrostis sp., Boerhaavia hispida, Cyperus compestris,</i>	1.0-1.5kg.	2.22-3.33	500-800	120-150
Flufenacet 60% DF					
Paddy (Transplanted)	<i>Echinochloa crusgalli Echinochloa colonum Cyperus iria</i>	120 gm	200 gm	500	90-110
Glufosinate Ammonium 13.5% SL (15% w/v)					
Tea	<i>Panicum repens, Borreria hispida, Imperata cylindrical, Digitaria sanguinalis, Commelina benghalensis, Ageratum conyzoides, Eleusine indica, Paspalum conjugatum</i>	0.375-0.500	2.5-3.3	375-500	15

Cotton	<i>Echinochloa sp.</i> <i>Cynodon dactylon</i> <i>Cyperus rotundus</i> <i>Digitaria marginata</i> <i>Dactyloctenium aegyptium</i>	375-450	2.5-3.0	500	96
Glyphosate 20.2% SL IPA salt					
Non Crop area	<i>Phyllanthus niruri</i> , <i>Ageratum conyzoides</i> , <i>Parthenium hysterophorus</i> , <i>Sorghum halepense</i> , <i>Amaranthus spinosus</i> , <i>Alternanthera sessilis</i> , <i>Cynodon dactylon</i> , <i>Cyperus rotundus</i> , <i>Echinochloa colonum</i> , <i>Trianthema portulacastrum</i>	0.82-1.23 kg	4.1-6.15	400-500	N/A
Glyphosate 41% SL IPA Salt					
Tea	<i>Arundinella bengalensis</i> <i>Axonopus compressus</i> <i>Cynodon dactylon</i> <i>Imperata cylindrical</i> <i>Kalm grass</i> <i>Paspalum scrobiculatum</i> <i>Polygonum perfoliatum</i>	0.820-1.230kg.	2.0-3.0	450	21
Non-cropped area	Sorghum halepense and other dicot & monocot weeds in general	0.820-1.230kg.	2.0-3.0	500	-
Glyphosate 54% SL (IPA Salt)					

Non Crop Area	<i>Ageratum conyzoides</i> <i>Alternanthera sessilis</i> <i>Commilina spp</i> <i>Cyperus spp</i> <i>Echinochloa sp.</i> <i>Eclipta alba</i> <i>Iscaemum rogosum</i> <i>Setaria spp</i>	1.8 kg	3.33 ltrs.	400-500	-
Glyphosate Ammonium Salt 5% SL					
Tea	<i>Ageratum conyzoides</i> <i>Biden pilosa</i> <i>Boreria latifolia</i> <i>Cynodon dactylon</i> <i>Cyperus rotundus</i> <i>Digitaria sanguinalis</i> <i>Euphorbia spp.</i> <i>Imperata cylendrica</i> <i>Paspalum conjugatum</i>	1.5 kg.	30 ltrs.	500	7 days
Non Crop area	<i>Cynodon dactylon</i> <i>Cyprus rotundus</i> <i>Digera arvensis</i> <i>Digitaria sanguinalis</i> <i>Eragrostis minor</i> <i>Euphorbia spp.</i> <i>Parthenium</i> <i>hysterophorus</i> <i>Tribulus terrestris</i> <i>Xantrhium stremerium</i>	2 kg.	40 ltrs.	500	-
Glyphosate 71% SG (Ammonium Salt)					

Tea & Non Crop area	<i>Acalypha indica</i> <i>Ageratum conyzoides</i> <i>Cychorium intybus</i> <i>Digera arvensis</i> <i>Cynodon dactylon</i> <i>Cyperus rotundus</i> <i>Digitaria sanguinalis</i> <i>Eragrostis spp.</i> <i>Ipomea digitaria</i> <i>Paspalum conjugatum</i> <i>Sida aculata</i>	2.13 kg	3.0 kg.	500	7
Halosulfuron Methyl 75% WG					
Sugarcane	Cyperus rotundus	60-67.5	80-90	375	294
Haloxypol R Methyl 10.5% w/w EC					
Soyabean	Brachiaria sp. Digitaria sanguinalis Dinebra arabica Echinochloa sp. Eleusine indica Eragrostis sp. Panicum isochmi	108-135	1000-1250	500	60
Imazethapyr Technical (H)					
Soyabean	Cyperus,diformis, Echinochloa colonum, E. crusgalli,Euphorbia hirta, Croton sperrisifeorus, Digera arvensis,	100	1000	500-600	75
Ground nut	Commelina bengalensis Cyperus difformis, Commelina benghalensis, Trianthema protulacasturm,Eragrot is pilosa	100-150	1000-1500	500-700	90
Imazethapyr 10% SL					

Soybean	<i>Cyperus difformis</i> <i>Echinochloa colonum</i> <i>E. crusgalli</i> <i>Euphorbia hirta</i> <i>Croton sperrisifeorus,</i> <i>Digera arvensis,</i> <i>Commelina</i> <i>Benghalensis</i>	100 gm	1.0ltr.	500-600	75
Groundnut	<i>Cyperus difformis</i> <i>Commelina</i> <i>benghalensis,</i> <i>Trianthema</i> <i>portulacasturm,</i> <i>Eragrostis pilosa</i>	100-150 gm	1.0-1.5 ltrs.	500-700	90
Isoproturon 50% WP					
Wheat	<i>Phalaris minor</i> <i>Avena fatua</i> <i>Poa annua</i>	1.0kg	2.0	750	-
Isoproturon 75% WP					
Wheat	<i>Phalaris minor</i> <i>Avena fatua</i> <i>Poa annua</i>	1.0kg	1.33 kg.	750	60 days
Linuron 50% WP					
Pea	<i>Anagallis arvensis,</i> <i>Chenopodium album,</i> <i>Chenopodium murale,</i> <i>Portulaca oleracea,</i> <i>Mielilotus indica,</i> <i>Melilotus alba,</i> <i>Medicago denticulata</i> <i>Fumeria parviflora,</i> <i>Echinochloa crusgalli,</i> <i>Poa annua.</i>	0.625-1.0 kg	1.25-2.0	500	80-90
MCPA, Amine salt 40% WSC					

Transplanted Rice	<i>Cyperus rotundus</i> <i>Impmoea reptans</i> <i>Ammania baccifera</i> <i>Lippia nodiflora</i> <i>Alternanthera sp.</i> <i>Ludwigia parviflora</i> <i>Marsilea quadrifoliata</i>	0.8-2.0 kg	2-5	400-600	
Wheat	<i>Chenopodium album</i> , <i>Asphodelus tenuifolius</i> <i>Fumaria parviflora</i> <i>Carthamus oxyacantha</i> <i>Launea sp.</i> , <i>Pluchia lanceolata</i> , <i>Melilotus indica</i> , <i>Vicia hirsuta</i> , <i>Lathyrus aphaca</i> , <i>Medicago denticulata</i> , <i>M. lupulina</i> , <i>Spergula arvensis</i> , <i>Argemone maxicana</i> , <i>Phyllathus niruri</i> .	1.0 kg	2.5	300-600	
Methabenzthiazuron 70% WP					
Wheat (PE –2DAS)	<i>Phalaris minor</i> , <i>Avena fatua</i> , <i>Avena ludoviciana</i> , <i>Poa annua</i> ,	1.05-1.4kg	1.5-2.0 kg.	700-1000	100
Wheat (Post –EM 30 DAS)	<i>Polypogon monspiliensis</i> , <i>Anagallis arvensis</i> , <i>Chenopodium album</i>	1.05-1.75kg	2.0-2.5 kg.	700-1000	100
Wheat (Early POE.16-18 DAS)	<i>Phalaris minor</i> , <i>Avena fatua</i> , <i>Avena ludoviciana</i> , <i>Chenopodium album</i>	0.7-0.87 kg	1.0-1.25 kg.	700-1000	100
Metolachlor 50% EC					

Soybean	<i>Echinochloa sp.</i> <i>Eleusine indica</i> <i>Digitaria sp.</i> <i>Panicum sp.</i> <i>Cyperus sp.</i> <i>Amaranthus viridis</i>	1.0 kg	2.0 ltrs.	600-750	-
Metribuzin 70% WP					
Soybean	<i>Digitaria spp.</i> <i>Cyperus esculentus</i> <i>Cyperus campestris</i> <i>Borreria spp.</i> <i>Eragrostis spp.</i>	0.35-0.525 kg	0.5-0.75kg.	750-1000	30
Wheat	<i>Phalaris minor</i> <i>Chenopodium album</i> <i>Melilotus spp.</i>	Medium soil- 0.175kg Heavy soil - 0.21kg	0.25 kg 0.30 kg.	500-750	120
Metsulfuron Methyl 20% WP					
Wheat	<i>Chenopodium album,</i> <i>Melilotus indica,</i> <i>Lathyrus aphaca,</i> <i>Anagallis arvensis,</i> <i>Vicia sativa,</i> <i>Cirsium arvense.</i>	4 gm	20 gm	500-600 + Surfactant (Iso-Octyl Phenoxy-Poloxethanol 12.5%)@ 500 ml/ha	80
Rice (transplanted)	<i>Cyperus rotundus,</i> <i>Spheanochlea spp.,</i> <i>Fimbristylis sp.</i> <i>Ludwigia parviflora</i> <i>Marsilea quadrifoliata</i>	4 gm.	20 gm.	500-600	60
Sugarcane	<i>Cyperus esculentus,</i> <i>Amaranthus viridis,</i> <i>Portulaca oleracea,</i> <i>Parthenium</i> <i>hysterophorus,</i> <i>Trianthema sp.,</i>	6	30	500-600 (Add non-ionic surfactant Iso-octyl-phenoxy-poloxethan	346

	<i>Cleome viscosa</i> , <i>Solanum sp.</i> , <i>Commelina benghalensis</i> , <i>Euphorbia sp.</i> , <i>Digeria sp.</i>			ol 12.5% @ 2ml per liter of spray volume (0.2%)	
Metsulfuron Methyl 20% WG					
Wheat	<i>Chenopodium album</i> <i>Melilotus indioca</i> <i>Melilotus alba</i> <i>Lathyrus aphaca</i> <i>Anagalis arvensis</i> <i>Vicia sativa</i> <i>Rumex denticulate</i> <i>Convolvulus arvensis</i> <i>Meedicago denticulate</i>	4 gm.	20 gm.	500-600 + Surfactan t (Iso- Octyl Phenoxy- Poloxeth anol 12.5%) @0.2%	76
Transplanted Rice	<i>Monochoria vaginalis</i> <i>Ludwigia parviflora</i> <i>Ludwigia adscendens</i> <i>Marselea quadrifoliata</i> <i>Eclipta alba</i> <i>Oxalis minima</i> <i>Dapatorium juncecum</i> <i>Commelina benghalensis</i> <i>Ammania baccifera</i> <i>Sphenoclea zeylanica</i> <i>Caesulia axillaries.</i>	4 gm	20 gm.	500-600 + Surfactan t (Iso- Octyl Phenoxy- Poloxeth anol 12.5%) @0.2%	71
Orthosulfamuron 50% WG					
Transplanted Rice (Paddy)	<i>Echinochloa spp.</i> (Barnyard grass) <i>Cyperus spp.</i> (Nut grass) <i>Scirpus spp.</i> <i>Ludwigia parviflora</i> (water crest) <i>Fimbristylis spp.</i> (Hoor grass) <i>Rotala spp.</i>	60-75	150 3 DAT	500	65
Oxadiargyl 80% WP					

Transplanted Rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> , <i>Cyperus iria</i> , <i>Cyperus</i> <i>difformis</i> , <i>Eclipta alba</i> <i>Ludwigia quadrifoliata</i>	100gm	0.125 kg.	500	97
Sunflower	<i>Echinochloa colona</i> <i>Dactyloctenium</i> <i>aegyptium</i>	240	300	500	81
Oxadiargyl 6%EC					
Transplanted Rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> ,	100gm	1.66 ltrs	500	97
Cumin	<i>Cyperus iria</i> , <i>Cyperus</i> <i>difformis</i> , <i>Eclipta alba</i> <i>Ludwigia quadrifoliata</i> <i>Chenopodium album</i> <i>Remex sp.</i> , <i>Melilotus indica</i> , <i>Asphodelus tenuifolius</i>	60-75gm	1.0-1.25 ltrs.	500	87
Mustard	<i>Chenopodium album</i> , <i>Melilotus sp</i>	90	1500	500	35
Oxadiazon 25% EC					
Transplanted Rice	<i>Echinochloa crusgalli</i> <i>E.colonum</i> <i>Cyperus iria</i> <i>C.difformis</i> <i>Marsilea</i> <i>quadrifoliata</i> , <i>Eclipta alba</i> , <i>Ludwigia sp.</i>	0.5kg	2.0 ltrs.	500	-
Oxyflourfen 0.35% GR					
Rice (Direct sown puddled or Transplanted)	<i>Echinochloa sp.</i> <i>Cyperus difformis</i> <i>Cyperus iria</i> <i>Eclipta alba</i> <i>Ludwigia parviflora</i> <i>Fimbristylis miliacia</i> , <i>Marsilea spp</i>	100-150 gm	30-40 kg.	-	-

Oxyflourfen 23.5% EC					
Rice (Direct sown as pre-emergence)	<i>Echinochloa sp.</i> <i>Cyperus iria, Eclipta alba,</i>	150-240 gm	650-1000	500	-
Tea	<i>Digitaria, Imperata, Paspalum, Borreria hispida,</i>	150-250 gm	650-1000	500-750	15 days
Onion	<i>Chenopodium album, Amaranthus viridis,</i>	100-200 gm	425-850	500-750	-
Potato	<i>Chenopodium, Coronpus Trianthema, Cyperus, Heliotropium</i>	100-200 gm	425-850	500-750	-
Groundnut	<i>Echinochloa colonum Digitaria arginata</i>	100-200 gm	425-850	500-750	-
Pendimethalin 30% EC					
Wheat	<i>Phalaris minor, Chenopodium album, Melilotus alba, Portulaca oleracea, Anagallis arvensis, Fumaria parviflora, Poa annua</i>	Light soil- 1.0 kg, Medium soil-1.25 kg, Heavy soil- 1.5 kg	3.3 ltr. 4.2 5.0	500-700 500-700 500-700	-
Rice (Transplanted & direct sown Upland)	<i>Echinochloa colona, E. crusgalli, Fimbristylis miliacea, Marselia quadrifoliata, Alternanthera sessilis, Ammonia baccifera, Ludwigra parviflora, Eclipta alba, Cyperus difformis</i>	Light to Heavy soil 1-1.5kg	3.3 – 5 Ltrs.	500-700	

Cotton	<i>Echinochloa spp.</i> <i>Euphorbia hirta</i> <i>Amarnanthus viridis</i> <i>Portulaca oleracea</i> <i>Trianthema spp.</i> <i>Eleusine indica</i>	0.75-1.25kg	2.5-4.165 ltrs	500-700	150
Soybean	<i>Echinochloa spp.</i> , <i>Euphorbia spp.</i> , <i>Amarnanthus viridis</i> , <i>Portulaca oleracea</i> , <i>Trianthema spp.</i> , <i>Eleusine indica</i>	0.75-1.0kg	2.5-3.3 ltrs.	500-700	110
Pigeon pea	<i>Digitaria sanguinalis</i> <i>Digitaria arvensis</i> <i>Amaranthus sp.</i> <i>Euphorbia hirta</i> <i>Trianthema sp.</i> <i>Cyperus sp.</i> <i>Eragrostis sp.</i>	0.7 – 1.00	2.5 – 3.33	500	133
Pendimethalin 5 % G					
Rice (Transplanted & Direct sown puddled)	<i>Echinochloa colona</i> , <i>E. crusgalli</i> , <i>Fimbristylis miliacea</i> , <i>Marselia quadrifoliata</i> , <i>Alternanthera sessilis</i> , <i>Ammonia baccifera</i> , <i>Ludwigra parviflora</i> , <i>Eclipta alba</i> , <i>Cyperus difformis</i>	1.0-1.5 kg	20-30 kg	-	-
Pendimethalin 38.7% CS					
Soybean	<i>Echinochloa colonum</i> <i>Dinebra arabuica</i> <i>Digitaria sanguinalis</i> <i>Bracharia mutica</i> <i>Dactyloctinum aegyptium</i> <i>Portulaca oleracea</i> <i>Amaranthus viridis</i> <i>Euphorbia geniculata</i> <i>Cleome viscose</i>	580.5-677.25gm	1500-1750	500	40

Cotton	Panicum repens, Digitaria sanguinalis, Brachiaria mutica (Grasses), Pennisetum purpureum, Cyperus rotundus (sedge), Lantana camjara, Portulaca oleracea, Eclipta prostrate, Commelina benghalensis (Broad leaves weeds)	677.27	1500- 1750	500	101
Chilli	Panicum repens, Digitaria sanguinalis, Elusine indica, Dinebra arabiaca, Echinochloa colonom, Portulaca oleracea, Commelina benghalensis, Aramthus blitum, Chenopodium album	677.27	1500- 1750	500	98
Onion	<i>Echinochloa colonom,</i> <i>Cyperus</i> <i>rotundus(Sedge)</i> <i>Cynodon dactylon</i> <i>Dinebra Arabic</i> <i>Euphor beageneculata</i> <i>Commelina</i> <i>bengalensis</i> <i>(Broad Leave weeds)</i>	580.50- 677.25g	1500- 1750	500	104
Pinoxaden 5.1% EC					
Wheat	<i>Phalaris minor</i> (Canary grass) <i>Avena ludoviciana</i> (Wild oat)	40-45 g	800-900 ml 30-35 DAS	225-300	90
Penoxsulam 21.7 % SC					

Rice (Transplanted)	Ammania bacifera, Cyperus difformis, Echinochloa colonum , Echinochloa crusgalli, Cyperus iria, Fimbristylis miliacea, Ludwigia spp. Monochoria spp. Sphenoclea zeylanica,	22.5 to 25 (pre- emergence 0-5 DAT) 20 to 22.5 (post- emergence 10-12 DAT)	93.7 to 104.2 83.3 to 93.7		60
Pretilachlor 37%EW					
Transplanted Rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis</i> <i>Cyperus iria</i> <i>Digitaria sanguinalis</i> <i>Fimbristylis miliacea</i> <i>Eclipta alba</i> <i>Ludwigia parviflora</i> <i>Monochoria vaginalis</i>	0.60-0.75 kg	1.5-1.875 ltrs.	500	90
Pretilachlor 30.7% EC					
(Direct seeded rice under puddled condition)	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis</i> <i>Cyperus iria</i>	0.45- 0.60kg.	1.5-2.0 ltr.	500	110
Pretilachlor 50% EC					

Transplanted Rice	<i>Echinochloa crusgalli</i> <i>Echinochloa colonum</i> <i>Cyperus difformis</i> <i>Cyperus iria</i> <i>Fimbristylis miliacea</i> <i>Eclipta alba</i> <i>Ludwigia parviflora</i> <i>Monochoria vaginalis</i> <i>Leptochloa chinensis</i> <i>Panicum repens</i>	0.50-0.75 kg.	1.0-1.5 ltrs.	500-700	75-90
Propaquizafop 10% EC					
Soybean	<i>Echinochloa colonum</i> , <i>Echinochloa crusgalli</i> , <i>Digiteria sanguinalis</i> , <i>Dactyloctenium egyptium</i> , <i>Eleusine indica</i>	50-75 g	500-750	500-750	21
Blackgram	<i>Echinochloa colonum</i> , <i>Echinochloa crusgalli</i> , <i>Digiteria sanguinalis</i> , <i>Dactyloctenium egyptium</i> , <i>Eleusine indica</i>	75-100 g	750-1000	500-750	21
Onion	<i>Echinochloa colonum</i> , <i>Digiteria sanguinalis</i> , <i>Dactyloctenium egyptium</i> , <i>Phalaris minor</i>	62.5	625	500	7
Paraquat dichloride 24% SL					
Tea (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Imperata</i> <i>Setaria sp.</i> , <i>Commelina benghalensis</i> , <i>Boerraria hispida</i> , <i>Paspalum conjugatum</i> ,	0.2-1.0 kg	0.8-4.25 ltr (For season long weed control, use 2.5-5.0 ltr for initial application. For subsequent repeat spot application use 1 litre)	200-400	Not Necessary (For season-long weed control, use 2.5 to 5 lit for initial application. For subsequent repeat spot application use 1 litre)

Potato (Post-emergence overall / inter-row application at 5-10 % emergence)	<i>Chenopodium sp.</i> <i>Angallis arvensis</i> <i>Trianthema monogyna</i> <i>Cyperus rotundus</i> <i>Fumeria parviflora</i>	0.5 kg	2.0 ltr.	500	100
Cotton (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Digera arvensis,</i> <i>Cyperus iria,</i> <i>Trianthema monogyna,</i> <i>Corchorus spp.,</i> <i>Leucas aspera,</i> <i>Euphorbia spp.</i>	0.3-0.5 kg	1.25-2.0	500	150-180
Rubber (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Digitaria sp., Eragrostis sp., Fimbristylis sp.</i>	0.3-0.6 kg	1.5-2.5	600	N.A.
Coffee	<i>Digitaria marginata,</i> <i>paspalum Conjugatum,</i> <i>Ageratum, Conyzides,</i> <i>Borreria hispida,</i> <i>Euphorbia hirta,</i> <i>Commelina benghalensis, Eleusine indica</i>	250	1.0	400	N.A.
Sugarcane	<i>Echinochloa crusgalli,</i> <i>Digiteria sanguinalis,</i> <i>Eleusine indica,</i> <i>Physalis minima,</i> <i>Portulaca oleracea.</i>	500	2.0	500	270
Sunflower	<i>Cyperus iria,</i> <i>Eleusine indica,</i> <i>Eragrostis sp.,</i> <i>Chenopoduim album,</i> <i>Trianthema monogyna</i>	400	1.6	500	120
Rice [pre-plant (minimum tillage) before sowing/transplanting for controlling standing weeds]	<i>Echinochloa crusgalli,</i> <i>Cyperus iria,</i> <i>Ageratum conyzides,</i> <i>Commelina benghalensis,</i> <i>Marsilea quadriofoliata,</i> <i>Brachiaria mutica</i>	0.3-0.8 kg	1.25-3.5	500	N.A.

Wheat [pre-plant (minimum tillage) before sowing]	<i>Grassy & Broad leaf weeds</i>	1.0 kg	4.25 ltrs	500	120-150
Maize [pre-plant (minimum tillage) before sowing]	<i>Cyperus rotundus, Commelina benghalensis, Trianthema monogyna, Amaranthus sp., Echinochloa sp</i>	0.2-0.5 kg	0.8-2.0 ltrs	500	90-120
Maize (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Cyperus iria, Cyperus rotundus, Commelina benghalensis Amaranthus sp. Echinochloa sp Trianthema monogyna</i>	0.2-0.5 kg	0.8-2.0 ltrs	500	90-120
Grapes (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Cyperus rotundus Cynodon dactylon Convolvulus sp. Portulaca sp. Tridax sp.</i>	0.5 kg.	2.0ltrs.	500	90
Apple (Post-emergence directed inter row application at 2-3 leaf stage of weeds)	<i>Rosa moschata Rosa eglantaria Rubus ellipticus</i>	0.75 kg	3.25 ltrs	700-1000	N.A.
Aquatic weed control					
Water ways Canals, Ponds Etc	<i>Eichhornia crassipes Hydrilla Typha latifolia</i>	1000 1000 1000-2000	4.25 4.25 4.25-8.5	600-1000 1000 600 600-1000	N.A
Pyrazosulfuron Ethyl 10% WP					
Transplanted Rice	<i>Cyperus Iria, Cyperus difformis, Fimbristylis miliacea, Monochoria vaginalis, Ludwigia parviflora</i>	10-15 g	100-150	500-600	95

Pyrethiobac Sodium 10% EC					
Cotton (Gossypium)	<i>Trianthema Spp</i> <i>Amaranthus Spp</i> <i>Chenopodium Spp</i> <i>Digera Spp</i> <i>Celosia argentia</i>	62.5-75 gm	625-750	500	160
Pyrozosaluron Ethyl 70% WDG					
Transplanted Rice	<i>Echinicloa spp</i> , <i>Cyparus rotundus</i> , <i>Ludwigia parviflora</i>	21g	-	-	43
Quizalofop-ethyl 5% EC					
Soybean	<i>Echinochloa crusgalli</i> <i>E. colomum</i> <i>Eragrostis sp.</i>	37.5-50 gm.	0.75-1.0	500-600	95
Cotton	<i>Echinolchloa crusgalli</i> , <i>Echinochloa colomum</i> , <i>Dinebra retroflexa</i> <i>Digiteria marginata</i>	50.5	1000	500	94
Groundnut	<i>Echinochloa colomum</i> , <i>Dinebra retroflexa</i> <i>Dactyloctenium sp.</i>	37.5-50.0	750-1000	500	89
Black gram	<i>Eleusine indica</i> , <i>Dactyloctenium aegyptium</i> , <i>Digitaria sanguinalis</i> , <i>Eragrostis sp.</i> , <i>Paspalidium sp.</i> , <i>Echinochloa sp.</i> , <i>Dinebra ratroflexa</i>	37.5-50.0	750-1000	500	52
Onion	<i>Digitaria sp.</i> , <i>Eleusine indicia</i> , <i>Dactyloctenium aegyptium</i> , <i>Eragrostis sp.</i> ,	37.5-50.0	750-1000	375-450	7
Quizalofop-ethyl 10% EC					
Soyabean	Love grass (<i>eragrostipilosa</i>), Crab grass (<i>digitaria</i>	375-45.0	375-450	300-500	69-103

	sanguinalis0/ wild finger/ Makra grass Viper grass, Barnyard grass, sanwa/Samel, Brown top millet				
Quizalofop –p-tefuryl 4.41% EC					
Soybean	<i>Echinochloa spp.</i> <i>Dinebra arabica</i> <i>Digitaria sanguinalis</i> <i>Cynodon dactylon</i> <i>Hemarthria compressa</i> <i>Eleusine indica</i>	30-40 gm	750-1000 ml	400	30
Sulfentrazone 39.6% w/w SC					
Soybean	<i>Acalypha sp.</i> <i>Commelina sp.</i> <i>Digera sp.</i> <i>Cyprus sp.</i> <i>Echinochloa sp.</i> <i>Brachiaria sp.</i> <i>Dinebra sp.</i>	360	750	500	88
Sulfosulfuron 75% WG					
Wheat	<i>Phalaris minor</i> <i>Chenopodium sp.</i> <i>Melilotus alba</i>	25 gm	33.3 gm	200-250 + Cationic surfactan t 1250ml/h a	110
Tembotrione 34.4% SC					
Maize	<i>Trianthema potulacastrum,</i> <i>Echinochloa sp.</i> <i>Bracheria sp.</i>	120g	286ml	500L	55
Triallate 50% EC					
Wheat	<i>Avena fatua</i>	1.25 kg	2.5 kg.	250-500	150
Triasulfuron 20% WS (H)					

WHEAT	Chenopodium album, Anagallis arvensis,Medilotus alba,Rumex spp, Medicago denticulata,Fumeria pomiflora, Cronopus didymus,Spergula arvensis and Malvela perviflora	20	100	500	81
-------	--	----	-----	-----	----

HERBICIDE COMBINATIONS

Anilofos 24% +2,4-D ethyl Ester 32% EC					
Transplanted rice	Echinochloa crusgalli Echinochloa colonum Ischaemum rugosum Fimbristylis miliacea	(0.24+ 0.32) to (0.36 + 0.48) kg	1-1.5 ltrs.	300	90
Bensulfuron methyl 0.6%+Pretilachlor 6% GR					
Transplanted Rice	Echinochloa crusgalli , Echinochloa colonum, ,Cynodon dactylon Cyperus iria , Cyperus difformis , Cyperus rotundus , Fimbristylis miliacea, Ludwigia parviflora,Marselia quadrifolia, Enhydra fluctuans,Sphenoclea zeylanica, Eclipta alba, Ammania baccifera.	60 + 600 gm	10 kg	N.A.	88
Carfentrazone ethyl 0.43% + Glyphosate 30.82% EW					
Tea	<i>Ageratum conyzoides</i> <i>Bidenspilosa</i> <i>Borreria sp.</i> <i>Crassocephalumcr</i> <i>epidioides</i> <i>Cynadon sp.</i> <i>Cyperous sp.</i> <i>Digitaria sp.</i> <i>Eleusine indica</i> <i>Mimosa sp.</i> <i>Mltracarpus villosus</i> <i>Oxalis sp.</i>	12.90 +924.60	3000	500	7
Non-cropped area	<i>Ageratum conyzoides</i> <i>Axonopus sp.</i> <i>Brachiaria sp.</i> <i>Commelina sp.</i> <i>Cynodon dactylon</i> <i>Cyperous sp.</i> <i>Digitaria sp.</i> <i>Eleusine indica</i> <i>Imperata cylendrica</i> <i>Lantana camera</i>	12.90 +924.60	3000	500	--

	<i>Parthenium sp.</i>				
Carfentrazone ethyl 20% + Sulfosulfuron 25% WG					
Wheat	Phalaris minor, avena ludoviciana Chenopodium album, ,Medilotus alba,Rumex spp,	20+25 +750 ml Surfactant	100	300	110
Clodinafop Propargyl 15% + Metsulfuron Methyl 1% WP					
Wheat	<i>Phalaris minor,</i> <i>Avena fatua,</i> <i>Chenopodium album,</i> <i>Melilotus sp.,</i> <i>Fumaria parviflora,</i> <i>Vicia sativa,</i> <i>Rumex sp.,</i> <i>Anagallis arvensis,</i> <i>Coronopus didymus,</i> <i>Lathyrus sp.,</i> <i>Convolvulus arvensis</i>	60+4	400	375 (Add 1250 ml surfacta nt at the time of sparying)	100
Clodinafop propargyl 9% + Metribuzin 20% W/W					
Wheat	<i>Phalaris mi</i> <i>Chenopodium album,</i> <i>Melilotus sp</i> <i>Vicia sativa,</i> <i>Rumex sp</i> <i>Medicago sp</i> <i>Cronopus didymus</i> <i>Dinebra vetroflexa</i>	54+120	600	300	120
Clomazone 20%+2,4-D EE 30% EC					
Transplanted Rice	Echinochloa colonum , Echinochloa crusgalli, Cyperus iria, Cyperus difformis,Eclipta alba,Leptochloa chinensis,Panicum repens,Fimbristylis miliacea,Marsilea quadrifoliata, Ludwigia parviflora.	0.250- 0.375 Kg	1.25 ltrs.	500	100-110

Fenoxaprop-p-ethyl 7.77% w/w + Metribuzin 13.6% w/w EC					
Wheat	<i>Phalaris minor</i> (Little seed canary grass) <i>Chenopodium album</i> (Lambs quarter) <i>Lathyrus aphaca</i> (Meadow Pea) <i>Rumes Sp.</i> (Golden dock) <i>Melilotus spp.</i> (Sweet clover) <i>Avena ludoviciana.</i>	100+175	1250	375	110
Fluazifop-p-butyl 11.1% w/w + Fomesafen 11.1% w/w SL					
Soybean	<i>Echinochloa colona</i> <i>Digitaria sp</i> <i>Eleusine indica</i> <i>Dactyloctenium aegyptium</i> <i>Brachiaria reptans</i> <i>Commelina benghalensis</i> <i>Digera arvensis</i> <i>Trianthema sp.</i> <i>Phyllanthus niruri</i> <i>Aclypha indica</i> <i>Dinebra arbica</i>	250	1000	500	71
Ground nut	<i>Echinochloa colona</i> <i>Digitaria sp.</i> <i>Eleusine indica</i> <i>Dactyloctenium aegyptium</i> <i>Commelina benghalensis</i> <i>Eluopus villosus</i> <i>Indigofera glandulosa</i> <i>Chloris barbata</i> <i>Trianthema sp.</i>	C250	1000	500	82

	<i>Digera arvensis</i> <i>Cleome viscosa</i> <i>Phyllanthus niruri</i> <i>Amaranthus viridis</i> <i>Cyperus sp.</i>				
Hexazinone 13.2% + Diuron 46.8 % WP					
Sugarcane	Enchinochloa colonum Dactyloctenium aegyptium Trianthema monogyna Amaranthus viridis Ipomea spp Cyperus rotundus Cyperus esculentus Setaria spp Parthenium hysterophorus Euphorbia hirta	1200 gm (264+936)	2 Kg	500	282-306
Imazamox 35% + Imazethapyr 35% WG					
Soybean	<i>Echinochloa sp.</i> <i>Dinebra arabica</i> <i>Digitaria sp.</i> <i>Brachiaria mutica</i> <i>Commelina benghalensis</i> <i>Euphorbia hirta</i>	70	100	375-500 Add surfactant (Cyspread) @ 1.5ml/litre of water + Ammonium sulphate @ 2.0 gm/litre of water	56
<i>Ground nut</i>	<i>Echinochloa colona</i> <i>Digitaria arvensis</i> <i>Commelina benghalensis</i> <i>Euphorbia hirta</i> <i>Amaranthus viridis</i> <i>Physalis minima</i>	70	100	375-500	83

Mesosulfuron Methyl 3% + Iodosulfuron Methyl Sodium 0.6% WG					
Wheat	Phalaris minor Medicago denticulata Chenopodium album Melilotus sp. Rumex sp. Anagallis arvensis Coronopus didymus Lathyrus aphaca Fumaria parviflora	(12+2.4 gm)	400 ml.	400-500 + Surfactant (Genopol LRO fluid) @ 500 ml/ha	96
Metsulfuron Methyl 10% + Chlorimuron ethyl 10% WP					
Transplanted Rice (Pre-emergence application-3 DAT)	Cyperus iria, Cyperus difformis, Fimbristylis miliaceae, Eclipta alba, Ludwigia parviflora, Cyanotis axillaries, Monocoria vaginalis, Marsilea quadrifoliata,	4gm	20 gm.	300	90
Metsulfuron Methyl 10% + Carfentrazone ethyl 40% DF					
Wheat	<i>Rumex sp.</i> <i>Medicago denticulate</i> <i>Malva parviflora</i> <i>Lathyrus aphaca</i> <i>Chenopodium album</i> <i>Melilotus alba</i> <i>Melilotus indica</i> <i>Anagallis arvensis</i> <i>Solanum nigrum</i> <i>Vicia sativa</i> <i>Convolvulus arvensis</i>	25	50	300	100
Oxyflurofen 2.5% + Glyphosate (Isopropyl amime salt)41% SC(w/w)					
Tea	Ageratum Conyzoids Cyperous sp Borreriabispida Pospalumcon jugatum Digitaria ciliaris	50+820	2000	500L/ha	14

Pendimethalin 30%+ Imazethapyr 2% EC					
Soybean	Echinochloa crusgalli Digera arvensis Commelina benghalensis, Amaranthus viridis Portulaca oleracea	(750+50) to (900+60) gm	2.5-3.0 ltrs	500-600	90
Pretilachlor 6% + pyrazosulfuron Ethyl 0.15%(H)					
Paddy	Grassy weeds, Broad Leave, Sedges	600+15	10	-	83
Propaquizafop 2.5% + Imazethapyr 3.75% w/w ME					
Soybean	<u>Grassy weeds:</u> Dactloctnium aegyptium Echinochloa acolonum Eleusine indica Digitaria sanguinalis <u>BLW:</u> Commelina Benghalensis Euphorbia hirta Digera arvensis Amaranthus viridis	50+75	2000	500	80
Sulfosulfuran 75%+ Metsulfuron Methyl 5%WG					
Wheat	Phalaris minor, Chenopodium sp., Medicago denticulata, Coronopos dedymus, Rumex spp. Melilotus alba, Anagallis arvensis	(30+2)	40 gm	250-500 + surfactant 1250 ml/ha	110
Sodium Aceflourofen 16.5% + Clodinafop Propargyl 8% EC					

Soybean	<i>Acalypha indica</i> , <i>aegyptium</i> , <i>Alternanthera v</i> <i>philoxeroides</i> , <i>Amaranthus spp.</i> , <i>Celosia argentea</i> , <i>Cleome viscosa</i> , <i>Commelina</i> <i>benghalensis</i> , <i>Dactyloctenium</i> <i>Digera arvensis</i> , <i>Digitaria sanguinalis</i> , <i>Echinochloa spp.</i> , <i>Eleusine indica</i> , <i>Euphorbia spp.</i> , <i>Parthenium spp.</i> , <i>Phyllanthus niruri</i> , <i>Physalis minima</i> , <i>Stellaria media</i> <i>Trianthema monogyna</i>	80 + 165	1000	500	61
---------	---	----------	------	-----	----